

THEORETICAL STUDY OF  $M^+—RG_2$  ( $M^+ = CA—RA$ ;  $RG = HE—RN$ )

ANNA ANDREJEVA, *School of Chemistry, University of Nottingham, Nottingham, United Kingdom*;  
ADRIAN GARDNER, *Department of Chemistry, Emory University, Atlanta, GA, USA*; JACK B GRANEK, *Center for Free-Electron Laser Science, Max Planck Institut für Kernphysik, Hamburg, Germany*;  
RICHARD PLOWRIGHT, *School of Chemistry, University of Nottingham, Nottingham, United Kingdom*;  
WILLIAM BRECKENRIDGE, *Department of Chemistry, The University of Utah, Salt Lake City, UT, USA*;  
TIM WRIGHT, *School of Chemistry, University of Nottingham, Nottingham, United Kingdom*.

*Ab initio* calculations were employed to investigate  $M^+—RG_2$  species. Trends in binding energies,  $D_e$ , bond lengths, and bond angles are discussed and rationalised by analysing the electronic wavefunction. Mulliken, natural population, and atoms-in-molecules (AIM) population analyses are performed. It is found that some complexes are linear whereas others are bent. Those results are discussed in terms of hybridization and the various interactions present in these species. The work is a continuation from a previously published study<sup>a</sup> where Group 2  $Be^+—RG_2$  and  $Mg^+—RG_2$  species are investigated and compared with Group 1  $Li^+—RG_2$  and  $Na^+—RG_2$  species.

---

<sup>a</sup>A. Andrejeva, A. M. Gardner, J. B. Graneek, R. J. Plowright, W. H. Breckenridge and T. G. Wright, *J. Phys. Chem. A.*, 117, 13578 (2013)